




# Animal

## The international journal of animal biosciences



### Review: Markers and proxies to monitor ruminal function and feed efficiency in young ruminants



L. Cheng<sup>a,\*</sup>, G. Cantalapiedra-Hijar<sup>b</sup>, S.J. Meale<sup>c</sup>, I. Rugoho<sup>d</sup>, A. Jonker<sup>e</sup>, M.A. Khan<sup>e</sup>, O. Al-Marashdeh<sup>f</sup>, R.J. Dewhurst<sup>g</sup>

<sup>a</sup> Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Dookie Campus, 3647 Victoria, Australia

<sup>b</sup> Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, F-63122 Saint-Genès-Champanelle, France

<sup>c</sup> School of Agriculture and Food Sciences, Faculty of Science, University of Queensland, Gatton, 4343 Queensland, Australia

<sup>d</sup> Lely Australia Pty Ltd, 84 Agar Drive, Truganina 3029, Victoria, Australia

<sup>e</sup> AgResearch Limited, Grasslands Research Centre, Private Bag 11008, Palmerston North 4410, New Zealand

<sup>f</sup> Faculty of Agriculture and Life Sciences, Lincoln University, P.O. Box 85084, Lincoln, New Zealand

<sup>g</sup> Scotland's Rural College, King's Buildings, West Mains Road, Edinburgh EH9 3JG, United Kingdom

#### ARTICLE INFO

##### Article history:

Received 19 June 2020

Revised 10 July 2021

Accepted 16 July 2021

##### Keywords:

body measures  
breath sulphide and methane  
faecal lipids  
nitrogen isotopes  
purine derivatives

#### ABSTRACT

Developing the rumen's capacity to utilise recalcitrant and low-value feed resources is important for ruminant production systems. Early-life nutrition and management practices have been shown to influence development of the rumen in young animals with long-term consequences on their performance. Therefore, there has been increasing interest to understand ruminal development and function in young ruminants to improve feed efficiency, health, welfare, and performance of both young and adult ruminants. However, due to the small size, rapid morphological changes and low initial microbial populations of the rumen, it is difficult to study ruminal function in young ruminants without major invasive approaches or slaughter studies. In this review, we discuss the usefulness of a range of proxies and markers to monitor ruminal function and nitrogen use efficiency (a major part of feed efficiency) in young ruminants. Breath sulphide and methane emissions showed the greatest potential as simple markers of a developing microbiota in young ruminants. However, there is only limited evidence for robust indicators of feed efficiency at this stage. The use of nitrogen isotopic discrimination based on plasma samples appeared to be the most promising proxy for feed efficiency in young ruminants. More research is needed to explore and refine potential proxies and markers to indicate ruminal function and feed efficiency in young ruminants, particularly for neonatal ruminants.

© 2021 The Authors. Published by Elsevier B.V. on behalf of The Animal Consortium. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

#### Implications

Simple measurements of sulphides and methane in breath could be used to provide a practical and non-invasive tool to monitor the developing microbiota of young ruminants. Plasma nitrogen isotopic discrimination is a promising proxy for feed efficiency in young ruminants and could be applied through a simple blood testing programme. However, the review indicated a lack of published international literature on the development of markers and proxies for ruminal function and feed efficiency in young ruminants, which would complement the much larger body of research on husbandry of young ruminants.

#### Importance of the rumen in digesting forages

The rumen and its microorganisms (bacteria, protozoa, fungi and archaea) facilitate the utilisation of substrates that are not available to mammalian enzymes (Van Soest and Van Soest, 1996; Liem et al., 2001), producing absorbable substrates for the host ruminant (Bergman, 1990). Physical breakdown during ingestion and rumination, as well as in the rumen, make feeds more accessible for microbial colonisation (Cheng et al., 1980; McAllister et al., 1994). The simple sugars formed are used by the microbes to produce volatile fatty acids (VFAs), mainly acetate (used for fatty acid synthesis), propionate (used for glucose synthesis), and butyrate which are largely used as energy sources in the ruminant body. Proportions of these VFAs in the rumen are influenced by microbial community composition (Henderson et al., 2015; Seshadri et al., 2018) and ruminal conditions. The ruminal

\* Corresponding author.

E-mail address: [long.cheng@unimelb.edu.au](mailto:long.cheng@unimelb.edu.au) (L. Cheng).

conditions are influenced by intake rate, dietary forage to concentrate ratio, and nature (e.g., degradation rate, molecular structure) of the diet. In general, forage dominated diets stimulate acetate formation whereas, concentrate dominated diets promote propionate formation. Feeds with higher levels of starch and protein levels promote propionate production, yet simple sugars and hemicellulose promote butyrate production, and cellulose promotes acetate production (Bannink et al., 2006).

Feed must be retained in the reticulo-rumen for long enough to allow the microorganisms to effectively ferment and break down plant fibre (Liem et al., 2001). Ruminants have a filter system between the reticulum and omasum to extend the ruminal retention time for slow fermenting neutral detergent fibre, which is a major component of forage (Van Soest, 1996). The rate and amount of microbial protein synthesis are determined by the availability of energy and protein in the rumen (Tedeschi et al., 2000). Carbohydrates are the main energy source for bacteria, although they can also be used as carbon skeletons for protein synthesis in combination with ammonia, amino acids or small peptides (Bach et al., 2005; Lanzas et al., 2008). Degradation of proteins yields peptides and amino acids, which are utilised by the microbes (transamination) or deaminated to yield VFAs, carbon dioxide and ammonia (Tamminga, 1979; Bach et al., 2005). The ammonia that exceeds the capacity of microbial growth is absorbed through the ruminal wall, converted into urea and circulated back into the rumen via saliva or excreted in the urine.

### Morphological development of the rumen in young ruminants

The development of the rumen and microbial colonisation is a two-way interaction between the host and microbial communities. Morphological development of the rumen is promoted by the consumption of solid feed. The associated production and absorption of VFAs as fermentation end-products stimulate the development of ruminal papillae, enabling their absorption and facilitating further epithelial metabolism (Sander et al., 1959; Suárez et al., 2006). Butyrate is the greatest stimulator of epithelial length and function, followed by propionate. Conversely, it is the physical structure of substrates like roughages, which expand ruminal volume, contribute to muscular development (Tamate et al., 1962; Stobo et al., 1966), and stimulate rumination and flow of saliva to the rumen (Hodgson, 1971).

The main enzymatic activities (fibrolysis, amylolysis, proteolysis, and ureolysis) of ruminal microbiota have been observed in the rumen from four (Sahoo et al., 2005) or ten (Kmet et al., 1986) days of age. Over 60 glycoside hydrolase microbial genes have been observed in the rumen during the early stages of life, suggesting great potential for plant carbohydrate metabolism even in the absence of regular plant cell wall intake (Li et al., 2012). As a calf grows, the ketogenic capacity of the rumen must develop to that of a mature rumen, as 60–80% of all VFAs are absorbed across the ruminal wall, with 75–90% of absorbed butyrate being metabolised by the ruminal epithelium (Allen, 1997).

### Microbial development in the rumen of young ruminants

Microbial inoculation of the rumen was considered to begin immediately after birth, through contact with the vaginal canal, faecal material, colostrum, skin and saliva of the dam. Yet recently, methanogens, fibrolytic bacteria, and Proteobacteria were detected in the rumen of calves less than 20 minutes after birth (Guzman et al., 2015). Quantification of bacterial and archaeal RNA (Malmuthuge et al., 2015) suggests that inoculation may in fact occur prior to birth, with rapid shifts occurring in the first days

of life as primo-colonising aerobic or facultative anaerobic bacteria shape the biotype for the strictly anaerobic microbes which sequentially establish thereafter (Jami et al., 2013). Similarities between establishment of the rumen and epimural microbial communities have been identified, as Proteobacteria were also found to be present at >90% of sequences from goat kids at birth (Rieu et al., 1990; Jiao et al., 2015; Wang et al., 2017). This is potentially due to their role in scavenging oxygen diffusing from the capillary network (Cheng et al., 1979), facilitating the establishment of anaerobic communities.

Recent studies (Li et al., 2012; Jami et al., 2013; Meale et al., 2016) suggest the preweaned rumen contains the same dominant phyla, Bacteroidetes, Firmicutes and Proteobacteria, as the more mature postweaned rumen, although relative abundance varies with age. Firmicutes increase after weaning (Jami et al., 2013; Meale et al., 2016; Meale et al., 2017a; 2017b), while Bacteroidetes, and specifically Prevotella, appear more dependent on solid food intake, than the removal of milk from the diet, reaching a stable abundance after 7 weeks of age (Meale et al., 2017a; 2017b), and once solid food consumption rises above 100 g per d, respectively (Rey et al., 2014; Meale et al., 2016; Meale et al., 2017a; 2017b). This indicates that the earlier a calf begins to consume solid feed, the sooner a ruminal bacterial community that is more representative of a mature ruminal develops. Furman et al. (2020) showed the effects of delivery method (spontaneous vs. caesarean) and diet, as well as random effects in early life on the development of the ruminal microbiome. Others (Roehe et al., 2016; Wallace et al., 2019) have identified host genetic effects on the ruminal microbiome. All of these genetic and early-life effects reinforce the need to quantify ruminal function in young ruminants – before, during and after weaning.

### Markers and proxies to monitor ruminal function in young ruminants

At birth, the rumen is sterile, and physically and metabolically underdeveloped. Initiation of solid feed consumption, acquisition of anaerobic microbes, the establishment of ruminal fermentation, growth of papillae, maturation of salivary function, and physical expansion of the rumen are achieved during the first ~ four months of life (Khan et al., 2011; Khan et al., 2016) in response to nutritional inputs and management practices (Yáñez-Ruiz et al., 2015; Steele et al., 2016; Meale et al., 2017a). For example, the provision of forage vs. concentrate to young ruminants provide a strong physical stimulus for the expansion of the rumen to increase its volume, physical development, and motility (Castells, 2013) and promote the development of rumination behaviour, saliva flow, and buffering capacity (Laarman and Oba, 2011; Khan et al., 2016). In summary, there is emerging evidence that early-life nutritional interventions influence ruminal development in young ruminants (Khan et al., 2016), with lifelong consequences on their welfare and performance (Khan et al., 2011; Soberon and Van Amburgh, 2013). There is renewed interest in finding mechanisms by which the host and diet can influence these developments, not least because there may be the longer-term setting of the interaction between the host, rumen and microbiome (Yáñez-Ruiz et al., 2015). However, there are serious constraints to the ability to use existing sampling techniques and limitations associated with the low ruminal volume and smaller microbial population. A number of approaches have been developed to study ruminal function less invasively and these have often been applied in studies with adult ruminants (see review by Dewhurst et al. (2000)). The purpose of the next section is to review these methods for prospects to be used with young ruminants; Table 1 provides a summary overview.

### Urinary purine derivatives

Urinary excretion of purine derivatives (**PDs**) has been used as an index of microbial protein supply in ruminants postweaning (Chen et al., 1990; Funaba et al., 1997). The basis of this approach is that nucleic acids leaving the rumen are of microbial origin (McAllan, 1980). Purines are important components of nucleic acids (the bases adenine and guanine) and absorbed purines are metabolised and excreted in urine as their end-products, which include allantoin, uric acid, xanthine, and hypoxanthine (Chen and Ørskov, 2004). Thus, urinary excretion of PD is quantitatively related to the mass of microbial protein supply to the host. However, this estimation is generally associated with error due to several factors including the endogenous contribution to urinary PD, variation in the purine to N ratio in the bacteria used as a reference (e.g., differences between solid- and liquid-associated bacteria (Bates et al., 1985)) and losses of PD through routes other than urinary excretion.

Firstly, saliva contains high levels of uric acid and allantoin, which are recycled to the rumen and degraded by the ruminal microbes (Chen et al., 1990; 1992) and estimates of microbial protein supply from urinary excretion of PD need to be corrected for such losses (Chen et al., 1992). However, because the development of salivary glands in calves varies and relates to time since weaning, there is a variable ratio of urinary PD to the intestinal flow of PD, and subsequent estimation of microbial protein (Funaba et al., 1997). Secondly, the endogenous PD contribution might also be variable during the period immediately after weaning. While endogenous PD production has been reported to be independent of the age (Funaba et al., 1997), Chen et al. (1992) showed that the abrupt removal of dietary protein supply increased endogenous allantoin production in sheep. More research is required to confirm the effect of endogenous production of PD on the accuracy of microbial protein prediction from urinary excretion of PD in young ruminants. Lastly, DMI increases with age in young ruminants after weaning, which means that a greater amount of feed PD may escape ruminal degradation and contribute to the total urinary excretion of PD (Shingfield, 2000), resulting in an overestimation of microbial protein. In conclusion, it appears that urinary excretion of PD is suitable to rank treatments based on relative differences in microbial protein synthesis, but not to give quantitative reference measurements for the individual animal, as suggested previously for adult ruminants (Shingfield, 2000).

### Faecal ether lipids (archaeol)

Recently, researchers have looked for distinctive components of methanogens (archaea) which could be measured directly in biological samples, either intact or following metabolism. The cell membranes of methanogenic archaea include unusual lipids, such as archaeol and caldarchaeol, which contain distinctive ether linkages. Archaeol is present in faeces from ruminants, but has not been detected in faeces from other herbivores (Gill et al., 2011). Therefore, faecal archaeol was proposed as a biomarker for methane (**CH<sub>4</sub>**) emissions from growing and adult ruminants. However, the relationship between CH<sub>4</sub> emissions and faecal archaeol concentrations is weak (Gill et al., 2011; McCartney et al., 2013; Schwarm et al., 2015), most likely due to differences in the passage rate of methanogens from the rumen (McCartney et al., 2014) suggesting that faecal archaeol has limited potential as a marker for methanogenesis. While the ability to distinguish ruminant faeces from non-ruminant faeces suggests that faecal archaeol might be a useful marker for the development and function of the rumen and methanogenesis, we are not aware of any published literature where such effects are measured during pre- and postweaning periods in young ruminants.

### Breath sulphide

Techniques to estimate the degradation of proteins by ruminal microbes have depended on the use of fistulated cows, so identifying markers of protein breakdown that can be accomplished in accessible samples is crucial to better understanding this aspect of ruminal function. Ruminal degradation of sulphur compounds, such as sulphates, methionine and cysteine, results in generation of hydrogen sulphide in the ruminal headspace gas (Dewhurst et al., 2007a). Some of the hydrogen sulphide is absorbed through the lungs (Dougherty et al., 1962) and subsequently metabolised to dimethyl sulphide, which is a distinctive component of cow's breath (Elliott-Martin et al., 1997). The multiple origins of sulphides in ruminal gases or breath make its measurement limited as a potential marker of protein degradation, but it does seem a viable option to monitor the establishment of the ruminal microbiota.

Much higher levels of hydrogen sulphide are generated when cattle consume water or feed contaminated with high levels of sulphate and this leads to a serious condition called polioencephalomalacia, which often results in death. A functioning ruminal microbial population is required for production of hydrogen sulphide and adaptation to high sulphate intakes (proliferation of sulphate-utilising microorganisms) can take several days (Lutnicki et al., 2014). Despite this adaptation, polioencephalomalacia was induced in 6-week old lambs offered a high sulphur diet (Gooneratne et al., 1989) demonstrating the potential to use sulphides as a marker for the activity of rumen microorganisms in young ruminants.

### Methane emissions

Anaerobic fermentation of feed in the rumen into acetate and butyrate also generates hydrogen (Janssen, 2010). This hydrogen is largely utilised by methanogens in the rumen, together with carbon dioxide, to form CH<sub>4</sub>, which is emitted by the ruminants. While hydrogenotrophic methanogenesis is the main pathway to CH<sub>4</sub> in the rumen, some diets can promote increased levels of other pathways, including from methyl-containing compounds (e.g., Neill et al., 1978). Both hydrogenotrophic and methylotrophic methanogenesis are microbial (archaeal) processes, so the CH<sub>4</sub> emitted is a quantitative proxy indicating ruminal fermentation. The DMI is the main driver of CH<sub>4</sub> production in postweaning growing cattle (Jiao et al., 2014; Jonker et al., 2016) and sheep (Muetzel and Clark, 2015) offered forage-based diets with CH<sub>4</sub> yields typically between 18 and 26 g/kg DMI. To our knowledge, there is little information available on CH<sub>4</sub> emissions from preweaned calves. One study with veal calves of over 15 weeks of age and 136 kg BW fed exclusively milk replacer found that negligible amounts of CH<sub>4</sub> were produced (<2 g/kg DMI) (Van den Borne et al., 2006). Dairy calves in two studies offered solid calf starter meal from three days of age (and total mixed ration with 50% hay from four weeks of age) already produced between 5 and 26 g CH<sub>4</sub>/kg DMI before weaning (when between two and eight weeks of age), while consuming 0.2–1.2 kg of solid feed (Muetzel, 2015). Up to week five of age, the CH<sub>4</sub> yield was lower (<16 g/kg DMI) than postweaning (21–26 g/kg DMI), and this appeared to be associated with low ruminal acetate/propionate ratio and high propionate concentration preweaning. Fermentation of feed into propionate leads to less hydrogen formation in the rumen and therefore less CH<sub>4</sub> formation (Van Nevel and Demeyer, 1996; Janssen, 2010). A meta-analysis indicated that the acetate/propionate ratio was higher in calves with access to forage preweaning and higher ruminal pH pre- and postweaning (Imani et al., 2017). The higher CH<sub>4</sub> yield in calves from week six to ten of age in the second study compared

**Table 1**

Markers/proxies to monitor rumen function in young ruminants.

Biomarker/proxy	Target use	Connection to physiology	Major factors affecting accuracy	Key references	Potential use
Urinary excretion of PD	Predicts rumen microbial (protein) synthesis	Nucleic acids leaving the rumen are essentially of microbial origin. Purines are important components of nucleic acids (the bases adenine and guanine) and absorbed purines are metabolised and excreted in urine as their end-products, which include allantoin, uric acid, xanthine and hypoxanthine.	1. Variable purine to nitrogen ratio in bacteria. 2. Losing PD through other excretion routes (e.g., uric acid and allantoin in saliva). 3. Endogenous PD contributions.	<a href="#">Bates et al. (1985)</a> ; <a href="#">Chen et al. (1992)</a> ; <a href="#">Gonzalez-Ronquillo et al. (2003)</a> ; <a href="#">Chen and Ørskov (2004)</a>	Yes
Faecal ether lipids (archaeol)	Predicts enteric CH <sub>4</sub> emissions	Unusual faecal lipids which originate from the membrane lipids of methanogens and so are related to methanogenesis.	1. Passage rate of methanogens from the rumen (selective retention). 2. Distribution and kinetics of methanogens in the rumen may contribute to genetic variation in CH <sub>4</sub> production.	<a href="#">Gill et al. (2011)</a> ; <a href="#">McCartney et al. (2013)</a> ; <a href="#">Schwarm et al. (2015)</a>	No
Breath sulphide	Indicates ruminal microbiota development	Rumen degradation of sulphur compounds generates hydrogen sulphide, which is absorbed and metabolised to sulphides, some of which are exhaled.	1. Water or feed contamination with sulphate. 2. Variable proportion excreted in urine. 3. A functioning rumen microbial population and adaptation is required to establish relevant microbes.	<a href="#">Raisbeck et al. (2008)</a> ; <a href="#">Cammack et al. (2010)</a> ; <a href="#">Lutnicki et al. (2014)</a>	Yes
CH <sub>4</sub> emissions	Indicates ruminal microbiota development	Anaerobic fermentation of feed in the rumen generates hydrogen, which is largely utilised by methanogens to form CH <sub>4</sub> .	1. Preweaning and young ruminants have low CH <sub>4</sub> emissions. 2. Different methods (e.g., laser CH <sub>4</sub> detector vs. chambers) to measure CH <sub>4</sub> have very different accuracies.	<a href="#">Chagunda (2013)</a> ; <a href="#">Muetzel (2015)</a>	Yes
OBCFA	Monitors rumen function, including microbial synthesis and volatile fatty acid proportions	OBCFAs are hardly found in feedstuffs but are present at higher levels in rumen microbial lipids; these appear in animal lipids, including blood and milk.	1. Limited studies in young animals. 2. Some postruminal synthesis or modification can affect OBCFA levels.	<a href="#">Westreicher-Kristen et al. (2020)</a>	No

PD = purine derivatives; CH<sub>4</sub> = methane; OBCFAs = odd- and branched-chain fatty acids.



to the first study by Muetzel and Clark (2015) could be due to higher forage intake, though this was not specified.

Most respiration chamber facilities are designed for work with larger growing and adult cattle and are often not suited for work with preweaning and young ruminants, which may have extremely low CH<sub>4</sub> emissions. However, it might be possible to operate these large chambers at a lower airflow rate to enable measurement of low CH<sub>4</sub> emissions (Muetzel, 2015), and there are also respiration chambers for small ruminants and facilities where chamber size is adjustable allowing quantification of CH<sub>4</sub> in young calves (e.g., Van den Borne et al., 2006). The laser CH<sub>4</sub> detector is highly sensitive to CH<sub>4</sub> concentrations in gas samples and has been used to estimate CH<sub>4</sub> emissions from cattle based on the frequency and CH<sub>4</sub> concentrations in eructed ruminal gases (Chagunda, 2013). Preliminary studies at Scotland's Rural College have demonstrated the use of the laser CH<sub>4</sub> detector to monitor the onset and development of a functioning ruminant in calves pre- and postweaning (Dewhurst; personal communication). Future studies are needed to validate if the laser CH<sub>4</sub> detector can provide useful CH<sub>4</sub> data, which in turn reflects ruminal function in young ruminants.

#### Fatty acids

Over the years, there has been an increasing interest in the use of odd- and branched-chain fatty acids (OBCFAs) as potential biomarkers to monitor ruminal function in mature ruminants (Vlaeminck et al., 2006a). This approach is also based on looking for components of ruminal microbes which are present (intact or as metabolites) in accessible samples; OBCFAs are hardly found in feedstuffs, but are present at higher levels in microbial lipids. For example, Kim et al. (2005) showed that OBCFA can be useful markers to study ruminal microbial colonisation, but the patterns of OBCFA did not identify the types of bacteria colonising herbage. Vlaeminck et al. (2005) noted that OBCFA can be used as markers for the duodenal flow of microbial matter in dairy cows, especially where feed intake data are not available. In addition, other studies (Vlaeminck et al., 2006b; Dewhurst et al., 2007a; 2007b; Bhagwat et al., 2012) evaluated the potential of OBCFA in milk to predict ruminal proportions of VFA and showed a strong relationship between milk OBCFA and molar proportions of individual VFA in the rumen. However, the past studies document the use of OBCFA as potential markers to monitor ruminal function used mature ruminants and many used milk samples. The potential of OBCFA to monitor ruminal function in young ruminants has not been explored.

#### Importance of feed efficiency for ruminant production

Feed is a major and variable input cost in ruminant production systems. Improvement in feed utilisation and conversion into products (feed efficiency; FE) is crucial, as it can lead to a substantial increase in productivity, profitability and potential gains in sustainability. While there are different ways to define FE depending on the production system, stock class and type of saleable products, most of the literature refers to FE as feed conversion efficiency (FCE), which is the product output per unit of feed intake, such as BW gain/DMI in growing sheep. Further, residual feed intake (RFI) defines the difference between actual and expected feed intake, based on BW and growth of the animal, and it measures FE that is independent of BW gain and mature body size (Crews, 2005). The RFI is increasingly used by animal breeders as a way to avoid selection for FE leading to correlated increases in animal size.

#### Challenges to quantify feed efficiency in young ruminants

There are many challenges when seeking to quantify FE in young ruminants, particularly in a grazing system or neonatal stage, where it is difficult to measure intake accurately. Ruminant FE measurements involve two components: product output (e.g., BW gain and milk production) and feed intake. The optimum test duration to accurately measure individual FE in growing ruminants ranges from 42 (Wang et al., 2006) to 100 days (Archer and Bergh, 2000). The International Committee for Animal Recording recommends a minimum period of 60 days, together with an adjustment period of at least 21 days, in which both individual animal feed intake and routine recording of animal BW are applied to remove as much of the non-genetic variation as possible. The recommended length for FE recording is a compromise between accuracy and minimum cost. In general, the duration for measuring an animal trait depends on its repeatability (i.e., time consistency or reliability), together with the frequency of the measurement (e.g., weekly vs. monthly). Repeatability or intra-class correlation coefficient is a measure of the tendency of animals to maintain their ranking over time and gives information about the magnitude of measurement errors (within-animal variance) compared to phenotypic variability (between-animal variance). Therefore, more repeatable animal traits subjected to less errors need less time to be accurately measured compared to those that are less repeatable.

In growing beef cattle, repeatability of DMI was reported to range between 0.51 and 0.70, whereas that for BW gain ranges between -0.03 and 0.21 (Kelly et al., 2010; Coyle et al., 2016). Thus, the measurement of DMI is not a critical trait determining the duration of the FE test (Archer et al., 1997). Repeatability of milk yield is higher than BW gain, ranging between 0.32 and 0.53 according to different estimates (7 studies summarised by Roman et al. (2000)). Thus, the time required to rank lactating cows according to their FE could be expected to be significantly shorter than the time required to rank growing ruminants. The exploration of markers and proxies of FE is needed to overcome the issues associated with the length and cost of measuring these traits, and will be essential in most field conditions where a reliable direct measurement of intake and performance is not possible or at least extremely challenging.

#### Markers and proxies to monitor feed efficiency

The use of markers and proxies to monitor FE has focused on growing or lactating animals, particularly the later stage of growing and finishing cattle. The next section reviews markers and proxies for prospects to be used with young ruminants (Table 2 provides a summary overview).

##### Body condition score and BW

The use of BW and BCS as proxies for FE is relatively simple, inexpensive and easy to implement on-farm (Negussie et al., 2017). Talebi (2012) showed a positive correlation ( $r^2 = 62\%$ ) between final BW and FCE in lambs. Similarly, several studies with young ruminants (Arthur et al., 1996; Basarab et al., 2003; Herd et al., 2016) noted a positive correlation between BCS and RFI, but the accuracy of the prediction varies. Other studies using young beef cattle reported a weak correlation between RFI and BW (Herd and Bishop, 2000; Schenkel et al., 2004). Similarly, studies have shown no relationship between RFI and BW in growing bulls (Arthur et al., 2001a; 2001b), steers (Nkrumah et al., 2004), beef heifers (Kelly et al., 2010) and growing dairy heifers (Green et al., 2013). Current literature shows that although there are

**Table 2**  
Markers/proxies to monitor feed efficiency in young ruminants.

Biomarker/proxy	Target use	Connection to physiology	Factors affecting accuracy	Key references	Potential use
BCS and BW gain	Indicates FE	BCS is related to body fat deposition, which is a major contributor to BW gain. BW gain contributes to the calculation of FE.	1. Age 2. Genetics 3. Nutrition 4. Measurement duration	<a href="#">Herd and Bishop (2000)</a> ; <a href="#">Schenkel et al. (2004)</a> ; <a href="#">Talebi (2012)</a>	BCS – No BW gain – may be useful for feed conversion efficiency, but not for residual feed intake
CH <sub>4</sub> emissions	Indicates FE	Methanogenesis is an energy loss during fermentation of feed in the rumen, thus CH <sub>4</sub> emissions contribute to inefficient use of feed.	1. Between-animal variation in energy losses from the same diets. 2. Difficulties to measure CH <sub>4</sub> emissions accurately.	<a href="#">Johnson and Johnson (1995)</a> ; <a href="#">Hegarty et al. (2007)</a>	May be
Blood- and milk-based markers	Indicates FE/nitrogen use efficiency	Biomarkers like insulin-like growth factor-1, aspartate aminotransferase, urea nitrogen, plasma nitrogen isotopic fractionation ( $\Delta^{15}\text{N}$ ) are related to either energy or protein metabolism or both.	1. Genetics 2. Nutrition 3. Protein turnover rate 4. Sampling time	<a href="#">Richardson and Herd (2004)</a> ; <a href="#">Huhtanen et al. (2015)</a> ; <a href="#">Cantalapiedra-Hijar et al. (2018a)</a>	$\Delta^{15}\text{N}$ – Yes Others – May be
Wool and hair-based marker	Indicates FE	Wool/hair $\Delta^{15}\text{N}$ is related to protein turnover and deamination and transamination in liver, which in turn reflects nitrogen use efficiency. Nitrogen use efficiency is a major component of FE.	1. Genetics 2. Nutrition 3. Protein turnover rate 4. Sampling techniques	<a href="#">Cheng et al. (2015)</a> ; <a href="#">Meale et al. (2017b)</a>	May be

CH<sub>4</sub> = methane; FE = feed efficiency; BCS = body condition score.

overall low to moderate relationships between RFI and BW, BW cannot be confidently used as a proxy for RFI in young ruminants. The use of both BW and BCS as proxies to predict FCE in young ruminants requires further investigation.

#### Methane emissions

Emissions of CH<sub>4</sub> are a loss of energy for the animal and reduced CH<sub>4</sub> emissions might therefore be associated with improved FE. However, the relationship between FE (RFI in most cases) and CH<sub>4</sub> yield has been inconsistent with relationships having been neutral ([Hegarty et al., 2007](#); [Waghorn and Hegarty, 2011](#); [Freetly and Brown-Brandl, 2013](#); [Alemu et al., 2017](#)), positive ([Nkrumah et al., 2006](#)) and negative ([Mercadante et al., 2015](#); [Herd et al., 2016](#); [McDonnell et al., 2016](#)) in postweaned growing cattle. While CH<sub>4</sub> is an important loss of energy that can range from 2 to 12% of gross energy (GE) intake ([Johnson and Johnson, 1995](#)), it is more usually in the range from 4 to 8% of GE intake with even less variation between animals offered the same diet. The use of CH<sub>4</sub> as a proxy for FE in young ruminants needs more study.

#### Blood- and milk-based markers

Since major mechanisms underlying the between-animal variability in FE are related to animal metabolism ([Cantalapiedra-Hijar et al., 2018b](#)), it can be argued that markers at the metabolic level may be more reliable than those at the digestive level to detect differences in FE across individuals. The potential of several hormones, such as leptin, insulin and insulin-like growth factor-1, aspartate aminotransferase and albumin as markers of between-animal variation in FE, has been proposed by several authors ([Johnson and Johnson, 1995](#); [Richardson and Herd, 2004](#); [Kelly et al., 2010](#)), though a recent review found inconsistent results ([Cantalapiedra-Hijar et al., 2018a; 2018b](#)). This could be due to interactions among plasma hormones and diet, physiological stage, age of animals or even the sampling procedures ([Cantalapiedra-](#)

[Hijar et al., 2018a; 2018b](#)), which precludes their use as reliable markers of FE at the individual animal level.

Recently, there has been an increasing interest in studying metabolites, proteins and genes potentially related to between-animal variability in FE. A recent study by [Duarte et al. \(2019\)](#) identified a common pathway related to branch-chain amino acid degradation through a meta-analysis of genome-wide association studies on RFI. Branch-chain amino acids have an important role in protein synthesis and turnover, energy-consuming metabolic processes, and their degradation can contribute to gluconeogenesis. [Cantalapiedra-Hijar et al. \(2018b\)](#) also identified protein turnover rate as one of the main determinants of animal variability in FE, and metabolites related to amino acid metabolism and protein turnover have already been proposed as indicators of FE ([Richardson and Herd, 2004](#); [Karisa et al., 2014](#); [Meale et al., 2017a; 2017b](#)).

A promising marker of FE is based on the  $^{15}\text{N}$  natural enrichment of animal proteins over the diet (nitrogen isotopic discrimination;  $\Delta^{15}\text{N}_{\text{animal-diet}}$ ). Nitrogen exists as two stable isotopes in nature: the more abundant light  $^{14}\text{N}$ , and the heavy  $^{15}\text{N}$ . The  $\Delta^{15}\text{N}_{\text{animal-diet}}$  originates from the isotopic selectivity of enzymes, leading to different  $^{15}\text{N}$  natural abundance between substrates and products during metabolic reactions ([Gannes et al., 1998](#)). Transaminase and deaminase in the animal liver are involved in major amino acid catabolism, and they have been suggested as key factors in the origin of  $\Delta^{15}\text{N}_{\text{animal-diet}}$  ([Macko et al., 1986](#)). Therefore, ruminants  $\Delta^{15}\text{N}$  biologically link to protein metabolism (e.g., protein use efficiency or nitrogen use efficiency) ([Cantalapiedra-Hijar et al., 2018a](#)) and indirectly link to FE, as nitrogen use efficiency (NUE) is a major component of FE ([Nasrollahi et al., 2020](#)).

This new biomarker seems to reflect changes in NUE or FCE across dietary conditions ([Cheng et al., 2013](#); [Cantalapiedra-Hijar et al., 2018b](#)), but also across individuals under the same diet and condition ([Wheadon et al., 2014](#); [Cantalapiedra-Hijar et al., 2018a](#)). More studies are warranted to explore the potential and

limitation of this new isotopic biomarker of between-animal variation in FE as well as to assess how heritable it is. Several studies have demonstrated the significant negative relationship between FCE and  $\Delta^{15}\text{N}_{\text{plasma-diet}}$ , and between FCE and  $\Delta^{15}\text{N}_{\text{blood-diet}}$  (Cantalapiedra-Hijar et al., 2018a), with the fact that it is driven by the protein component of growth (i.e., NUE) suggested by the improvement in relationships when ultrasound-based estimates of body composition, such as fat deposition, are included in the relationship (Meale et al., 2018). However, it is unclear how this pertains to neonatal ruminants.

It is worth noting that although urea nitrogen content of blood or milk has also been proposed as a moderately heritable marker of NUE, it may not be able to capture the animal variability properly in NUE (Huhtanen et al., 2015). Furthermore, several studies have also proposed blood urea nitrogen as a marker for RFI, but its inconsistency across time (Richardson and Herd, 2004) precludes its use as a universal marker of FE. The fact that blood or milk urea nitrogen describes digestive rather than metabolic variations in the NUE (Hof et al., 1997) could partly explain why it fails to reflect the between-animal variability in NUE or FE.

#### Wool and hair-based marker

While  $\Delta^{15}\text{N}_{\text{plasma-diet}}$  showed its potential use as a biomarker to indicate FE in both large and small ruminants, a recent study showed that  $\Delta^{15}\text{N}_{\text{wool-diet}}$  was negatively correlated with FCE of growing sheep (Cheng et al., 2015). Though the study is a preliminary analysis, it highlights the potential to use a marker from non-invasive and easy to obtain samples, such as wool, to predict FE. Wool is known to provide a cumulative  $^{15}\text{N}$  signature, which may be used to indicate a cumulative change in FE over a longer period of time than blood. However, it is unclear how this pertains to neonatal ruminants. Further, hair from cattle demonstrated a similar potential to indicate FE (Meale et al., 2017b).

#### Bringing ruminal function and feed efficiency together, and what comes next?

Unifying the two areas of this review, there is a current interest in the relationship between ruminal function and FE in the context of selection for increased FE, as well as the possibility to manipulate it through persistent effects of early-life interventions. Richardson and Herd (2004) analysed the FE trait in beef cattle and suggested that digestion contributes a relatively small proportion of the trait in comparison with metabolic effects. At the same time, the ruminal metagenomics work of Roehe et al. (2016) suggests that ruminal processes have a strong relationship with FE. Clearly, there is still a lot to learn about the interactions between the host and microbiome in the rumen. Given the interest in genetic effects on FE and the long-term effects of early-life development of ruminal function, there is a real need to make measurements of ruminal function and FE on larger numbers of animals (genetic studies) offered with different diets (diet studies). Detailed simultaneous analysis of microbiomes and metabolomes in such large studies will help identify new markers or proxies that can be used to optimise both ruminal function and host FE.

#### Conclusion

The use of breath sulphide and  $\text{CH}_4$  emissions shows potential as non-invasive markers for the establishment of the rumen microbiota in young ruminants; however, their use as robust indicators of FE is limited at this stage. The use of  $^{15}\text{N}$  discrimination from plasma samples appears the most promising proxy for FE, which maintains its strong negative relationship with NUE and FCE across

varying animal ages, indicating its potential as a marker to facilitate larger scale studies with growing ruminants. More research is needed to explore potential proxies and markers to indicate ruminal function and FE in young ruminants, as the current available indicators are mostly developed for mature ruminants.

#### Ethics approval

Not applicable.

#### Data and model availability statement

Data mentioned in this publication can be found from cited literature.

#### Author ORCIDs

L. Cheng: <https://orcid.org/0000-0002-8483-0495>

G. Cantalapiedra-Hijar: <https://orcid.org/0000-0001-9486-8238>

S.J. Meale: <https://orcid.org/0000-0002-8417-1186>

I. Rugoho: <https://orcid.org/0000-0002-3792-3584>

A. Jonker: <https://orcid.org/0000-0002-6756-8616>

M.A. Khan: <https://orcid.org/0000-0001-7590-5128>

O. Al-Marashdeh: <https://orcid.org/0000-0001-6493-0395>

R.J. Dewhurst: <https://orcid.org/0000-0002-9357-7372>

#### Author contributions

LC and RD conceptualized and designed the review, LC, GC, SM, IR, AJ, MK, OA and RJ drafted different sections according to their expertise and revised the article.

#### Declaration of interest

No competing interests to report.

#### Acknowledgement

Authors want to acknowledge Md Safiqur Rahaman Shishir (The University of Melbourne, Australia), Mr. Zelin Li (The University of Melbourne, Australia) and Prof. Jianjun Hu (Tarim University, China), for providing useful discussion points and formatting this paper.

#### Financial support statement

The research received no specific grant from any funding agency, commercial or not-for-profit section.

#### References

- Alemu, A., Vyas, D., Manafiazar, G., Basarab, J., Beauchemin, K., 2017. Enteric methane emissions from low- and high-residual feed intake beef heifers measured using GreenFeed and respiration chamber techniques. *Journal of Animal Science* 95, 3727–3737.
- Allen, M.S., 1997. Relationship between fermentation acid production in the rumen and the requirement for physically effective fiber. *Journal of Dairy Science* 80, 1447–1462.
- Archer, J., Arthur, P., Herd, R., Parnell, P., Pitchford, W., 1997. Optimum postweaning test for measurement of growth rate, feed intake, and feed efficiency in British breed cattle. *Journal of Animal Science* 75, 2024–2032.
- Archer, J., Bergh, L., 2000. Duration of performance tests for growth rate, feed intake and feed efficiency in four biological types of beef cattle. *Livestock Production Science* 65, 47–55.
- Arthur, P., Archer, J., Johnston, D., Herd, R., Richardson, E., Parnell, P., 2001a. Genetic and phenotypic variance and covariance components for feed intake, feed



- efficiency, and other postweaning traits in Angus cattle. *Journal of Animal Science* 79, 2805–2811.
- Arthur, P.F., Herd, R.M., Wright, J., Xu, G., Dibley, K., Richardson, E.C., 1996. Net feed conversion efficiency and its relationship with other traits in beef cattle. *Proceedings of the Australian Society of Animal Production* 21, 107–110.
- Arthur, P., Renand, G., Krauss, D., 2001b. Genetic and phenotypic relationships among different measures of growth and feed efficiency in young Charolais bulls. *Livestock Production Science* 68, 131–139.
- Bach, A., Calsamiglia, S., Stern, M.D., 2005. Nitrogen metabolism in the rumen. *Journal of Dairy Science* 88, E9–E21.
- Bannink, A., Kogut, J., Dijkstra, J., France, J., Kebreab, E., Van Vuuren, A., Tamminga, S., 2006. Estimation of the stoichiometry of volatile fatty acid production in the rumen of lactating cows. *Journal of Theoretical Biology* 238, 36–51.
- Basarab, J., Price, M., Aalhus, J., Okine, E., Snelling, W., Lyle, K., 2003. Residual feed intake and body composition in young growing cattle. *Canadian Journal of Animal Science* 83, 189–204.
- Bates, D.B., Gillett, J.A., Barao, S.A., Bergen, W.G., 1985. The effect of specific growth rate and stage of growth on nucleic acid-protein values of pure cultures and mixed ruminal bacteria. *Journal of Animal Science* 61, 713–724.
- Bergman, E., 1990. Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiological Reviews* 70, 567–590.
- Bhagwat, A., De Baets, B., Steen, A., Vlaeminck, B., Fievez, V., 2012. Prediction of ruminal volatile fatty acid proportions of lactating dairy cows based on milk odd- and branched-chain fatty acid profiles: New models, better predictions. *Journal of Dairy Science* 95, 3926–3937.
- Cammack, K.M., Wright, C.L., Austin, K., Johnson, P., Cockrum, R., Kessler, K., Olson, K., 2010. Effects of high-sulfur water and clinoptilolite on health and growth performance of steers fed forage-based diets. *Journal of Animal Science* 88, 1777–1785.
- Cantalapiedra-Hijar, G., Abo-Ismael, M., Carstens, G., Guan, L., Hegarty, R., Kenny, D., McGee, M., Plastow, G., Relling, A., Ortigues-Marty, I., 2018a. Biological determinants of between-animal variation in feed efficiency of growing beef cattle. *Animal* 12 (suppl. 2), 321–335.
- Cantalapiedra-Hijar, G., Dewhurst, R., Cheng, L., Cabrita, A., Fonseca, A., Nozière, P., Makowski, D., Fouillet, H., Ortigues-Marty, I., 2018b. Nitrogen isotopic fractionation as a biomarker for nitrogen use efficiency in ruminants: a meta-analysis. *Animal* 12, 1827–1837.
- Castells, D., 2013. Effect of different fiber sources on the digestive function and development of calves 1 recurs electrònic, ISBN 9788449037818. PhD Thesis. The Autonomous University of Barcelona, Barcelona, Spain <https://ddd.uab.cat/record/126826>.
- Chagunda, M., 2013. Opportunities and challenges in the use of the Laser Methane Detector to monitor enteric methane emissions from ruminants. *Animal* 7 (suppl. 2), 394–400.
- Chen, X., Hovell, F.D., Ørskov, E., 1990. Excretion of purine derivatives by ruminants: recycling of allantoin into the rumen via saliva and its fate in the gut. *British Journal of Nutrition* 63, 197–205.
- Chen, X.B., Ørskov, E.R., 2004. Research on urinary excretion of purine derivatives in ruminants: past, present and future. In: Makkar, H.P.S., Chen, X.B. (Eds.), *Estimation of microbial protein supply in ruminants using urinary purine derivatives*. Springer, Dordrecht, The Netherlands, pp. 180–210. [https://doi.org/10.1007/978-1-4020-2844-1\\_21](https://doi.org/10.1007/978-1-4020-2844-1_21).
- Chen, X.B., Chowdhury, S.A., Hovell, F.D.D., Ørskov, E.R., Kyle, D.J., 1992. Endogenous allantoin excretion in response to changes in protein supply in sheep. *The Journal of Nutrition* 122, 2226–2232.
- Cheng, K.J., Fay, J.P., Howarth, R.E., Costerton, J.W., 1980. Sequence of events in the digestion of fresh legume leaves by rumen bacteria. *Applied and Environmental Microbiology* 40, 613–625.
- Cheng, K., McCowan, R., Costerton, J., 1979. Adherent epithelial bacteria in ruminants and their roles in digestive tract function. *The American Journal of Clinical Nutrition* 32, 139–148.
- Cheng, L., Logan, C.M., Dewhurst, R.J., Hodge, S., Zhou, H., Edwards, G., 2015. Effects of feed intake and genetics on tissue nitrogen-15 enrichment and feed conversion efficiency in sheep. *Journal of Animal Science* 93, 5849–5855.
- Cheng, L., Sheahan, A., Gibbs, S., Rius, A., Kay, J., Meier, S., Edwards, G., Dewhurst, R., Roche, J., 2013. Nitrogen isotopic fractionation can be used to predict nitrogen-use efficiency in dairy cows fed temperate pasture. *Journal of Animal Science* 91, 5785–5788.
- Coyle, S., Fitzsimons, C., Kenny, D., Kelly, A., McGee, M., 2016. Repeatability of feed efficiency in steers offered a high-concentrate diet. *Journal of Animal Science* 94, 719.
- Crews, J.D., 2005. Genetics of efficient feed utilization and national cattle evaluation: a review. *Genetics and Molecular Research: GMR* 4, 152–165.
- Dewhurst, R., Davies, D., Merry, R., 2000. Microbial protein supply from the rumen. *Animal Feed Science and Technology* 85, 1–21.
- Dewhurst, R., Kim, E., Evans, R., Cabrita, A., Fonseca, A., 2007a. Effects of dietary sulphur sources on concentrations of hydrogen sulphide in the rumen head-space gas of dairy cows. *Animal* 1, 531–535.
- Dewhurst, R., Moorby, J., Vlaeminck, B., Fievez, V., 2007b. Apparent recovery of duodenal odd- and branched-chain fatty acids in milk of dairy cows. *Journal of Dairy Science* 90, 1775–1780.
- Dougherty, R., Stewart, W., Nold, M., Lindahl, I., Mullenax, C., Leek, B., 1962. Pulmonary absorption of eructated gas in ruminants. *American Journal of Veterinary Research* 23, 205–212.
- Duarte, D., Newbold, C.J., Detmann, E., Silva, F., Freitas, P., Veroneze, R., Duarte, M., 2019. Genome-wide association studies pathway-based meta-analysis for residual feed intake in beef cattle. *Animal Genetics* 50, 150–153.
- Elliott-Martin, R., Mottram, T., Gardner, J., Hobbs, P., Bartlett, P., 1997. Preliminary investigation of breath sampling as a monitor of health in dairy cattle. *Journal of Agricultural Engineering Research* 67, 267–275.
- Freetly, H., Brown-Brandl, T., 2013. Enteric methane production from beef cattle that vary in feed efficiency. *Journal of Animal Science* 91, 4826–4831.
- Funaba, M., Kagiya, K., Iriki, T., Abe, M., 1997. Duodenal flow of microbial nitrogen estimated from urinary excretion of purine derivatives in calves after early weaning. *Journal of Animal Science* 75, 1965–1973.
- Furman, O., Shenhav, L., Sasson, G., Kokou, F., Honig, H., Jacoby, S., Hertz, T., Cordero, O.X., Halperin, E., Mizrahi, I., 2020. Stochasticity constrained by deterministic effects of diet and age drive rumen microbiome assembly dynamics. *Nature Communications* 11, 1–13.
- Gannes, L.Z., Del Rio, C.M., Koch, P., 1998. Natural abundance variations in stable isotopes and their potential uses in animal physiological ecology. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 119, 725–737.
- Gill, F., Dewhurst, R., Evershed, R., McGeough, E., O'Kiely, P., Pancost, R., Bull, I., 2011. Analysis of archaeal ether lipids in bovine faeces. *Animal Feed Science and Technology* 166, 87–92.
- Gonzalez-Ronquillo, M., Balcells, J., Guada, J., Vicente, F., 2003. Purine derivative excretion in dairy cows: Endogenous excretion and the effect of exogenous nucleic acid supply. *Journal of Dairy Science* 86, 1282–1291.
- Gooneratne, S., Olkowski, A., Christensen, D., 1989. Sulfur-induced polioencephalomalacia in sheep: some biochemical changes. *Canadian Journal of Veterinary Research* 53, 462.
- Green, T., Jago, J., Macdonald, K., Waghorn, G., 2013. Relationships between residual feed intake, average daily gain, and feeding behavior in growing dairy heifers. *Journal of Dairy Science* 96, 3098–3107.
- Guzman, C.E., Bereza-Malcolm, L.T., De Groef, B., Franks, A.E., 2015. Presence of selected methanogens, fibrolytic bacteria, and proteobacteria in the gastrointestinal tract of neonatal dairy calves from birth to 72 hours. *PLOS One* 10, e0133048. <https://doi.org/10.1371/journal.pone.0133048>.
- Hegarty, R., Goopy, J.P., Herd, R., McCorkell, B., 2007. Cattle selected for lower residual feed intake have reduced daily methane production. *Journal of Animal Science* 85, 1479–1486.
- Henderson, G., Cox, F., Ganesh, S., Jonker, A., Young, W., Abecia, L., Angarita, E., Aravena, P., Arenas, G.N., Ariza, C., 2015. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Scientific Reports* 5, 14567.
- Herd, R., Bishop, S., 2000. Genetic variation in residual feed intake and its association with other production traits in British Hereford cattle. *Livestock Production Science* 63, 111–119.
- Herd, R.M., Velazco, J., Arthur, P., Hegarty, R., 2016. Associations among methane emission traits measured in the feedlot and in respiration chambers in Angus cattle bred to vary in feed efficiency. *Journal of Animal Science* 94, 4882–4891.
- Hodgson, J., 1971. The development of solid food intake in calves 4. The effect of the addition of material to the rumen, or its removal from the rumen, on voluntary food intake. *Animal Science* 13, 581–592.
- Hof, G., Vervoor, M., Lenaers, P., Tamminga, S., 1997. Milk urea nitrogen as a tool to monitor the protein nutrition of dairy cows. *Journal of Dairy Science* 80, 3333–3340.
- Huhtanen, P., Cabezas-Garcia, E., Krizsan, S., Shingfield, K., 2015. Evaluation of between-cow variation in milk urea and rumen ammonia nitrogen concentrations and the association with nitrogen utilization and diet digestibility in lactating cows. *Journal of Dairy Science* 98, 3182–3196.
- Imani, M., Mirzaei, M., Baghbanzadeh-Nobari, B., Ghaffari, M., 2017. Effects of forage provision to dairy calves on growth performance and rumen fermentation: A meta-analysis and meta-regression. *Journal of Dairy Science* 100, 1136–1150.
- Jami, E., Israel, A., Kotser, A., Mizrahi, I., 2013. Exploring the bovine rumen bacterial community from birth to adulthood. *The ISME Journal* 7, 1069–1079.
- Janssen, P.H., 2010. Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Animal Feed Science and Technology* 160, 1–22.
- Jiao, H., Yan, T., Wills, D.A., Carson, A.F., McDowell, D.A., 2014. Development of prediction models for quantification of total methane emission from enteric fermentation of young Holstein cattle at various ages. *Agriculture, Ecosystems & Environment* 183, 160–166.
- Jiao, J., Huang, J., Zhou, C., Tan, Z., 2015. Taxonomic identification of ruminal epithelial bacterial diversity during rumen development in goats. *Applied and Environmental Microbiology* 81, 3502–3509.
- Johnson, K.A., Johnson, D.E., 1995. Methane emissions from cattle. *Journal of Animal Science* 73, 2483–2492.
- Jonker, A., Muetzel, S., Molano, G., Pacheco, D., 2016. Effect of fresh pasture forage quality, feeding level and supplementation on methane emissions from growing beef cattle. *Animal Production Science* 56, 1714–1721.
- Karisa, B., Thomson, J., Wang, Z., Li, C., Montanholi, Y., Miller, S., Moore, S., Plastow, G., 2014. Plasma metabolites associated with residual feed intake and other productivity performance traits in beef cattle. *Livestock Science* 165, 200–211.
- Kelly, A., McGee, M., Crews Jr., D., Fahey, A., Wylie, A., Kenny, D., 2010. Effect of divergence in residual feed intake on feeding behavior, blood metabolic



- variables, and body composition traits in growing beef heifers. *Journal of Animal Science* 88, 109–123.
- Khan, M., Bach, A., Weary, D., Von Keyserlingk, M., 2016. Invited review: Transitioning from milk to solid feed in dairy heifers. *Journal of Dairy Science* 99, 885–902.
- Khan, M., Weary, D., Von Keyserlingk, M., 2011. Invited review: Effects of milk ration on solid feed intake, weaning, and performance in dairy heifers. *Journal of Dairy Science* 94, 1071–1081.
- Kim, E.J., Sanderson, R., Dhanoa, M.S., Dewhurst, R.J., 2005. Fatty acid profiles associated with microbial colonization of freshly ingested grass and rumen biohydrogenation. *Journal of Dairy Science* 88, 3220–3230.
- Kmet, V., K. Boda, P. Javorský, and R. Nemcová, 1986. The enzymatic activity of rumen microflora in calves. *Journal of Animal Physiology and Animal Nutrition* 56, 73–77.
- Laarman, A., Oba, M., 2011. Effect of calf starter on rumen pH of Holstein dairy calves at weaning. *Journal of Dairy Science* 94, 5661–5664.
- Lanzas, C., Broderick, G., Fox, D., 2008. Improved feed protein fractionation schemes for formulating rations with the Cornell Net Carbohydrate and Protein System. *Journal of Dairy Science* 91, 4881–4891.
- Li, R.W., Connor, E.E., Li, C., Baldwin, V., Ransom, L., Sparks, M.E., 2012. Characterization of the rumen microbiota of pre-ruminant calves using metagenomic tools. *Environmental Microbiology* 14, 129–139.
- Liem, K.F., Bemis, W.E., Walker, W.F., Grande, L., Walker Jr., W.F., The University of California, 2001. Functional anatomy of the vertebrates: an evolutionary perspective, Volume 1, Edition 3. Harcourt College Publishers, San Diego, CA, USA.
- Lutnicki, K., Madej, E., Riha, T., Kurek, L., 2014. Polioencephalomalacia in ruminants caused by excessive amount of sulphur—a review. *Bulletin of the Veterinary Institute in Pulawy* 58, 321–326.
- Macko, S.A., Estep, M.L.F., Engel, M.H., Hare, P., 1986. Kinetic fractionation of stable nitrogen isotopes during amino acid transamination. *Geochimica et Cosmochimica Acta* 50, 2143–2146.
- Malmuthuge, N., Griebel, P.J., Guan, L.L., 2015. The gut microbiome and its potential role in the development and function of newborn calf gastrointestinal tract. *Frontiers in Veterinary Science* 2, 36.
- McAllan, A., 1980. The degradation of nucleic acids in, and the removal of breakdown products from the small intestines of steers. *British Journal of Nutrition* 44, 99–112.
- McAllister, T., Bae, H., Jones, G., Cheng, K.-J., 1994. Microbial attachment and feed digestion in the rumen. *Journal of Animal Science* 72, 3004–3018.
- McCartney, C., Bull, I., Dewhurst, R.J., 2014. Using archaeol to investigate the location of methanogens in the ruminant digestive tract. *Livestock Science* 164, 39–45.
- McCartney, C., Bull, I., Yan, T., Dewhurst, R.J., 2013. Assessment of archaeol as a molecular proxy for methane production in cattle. *Journal of Dairy Science* 96, 1211–1217.
- McDonnell, R., Hart, K., Boland, T., Kelly, A., McGee, M., Kenny, D., 2016. Effect of divergence in phenotypic residual feed intake on methane emissions, ruminal fermentation, and apparent whole-trait digestibility of beef heifers across three contrasting diets. *Journal of Animal Science* 94, 1179–1193.
- Meale, S.J., Auffret, M.D., Watson, M., Morgavi, D.P., Cantalapiedra-Hijar, G., Duthie, C.-A., Roehe, R., Dewhurst, R.J., 2018. Fat accretion measurements strengthen the relationship between feed conversion efficiency and nitrogen isotopic discrimination while rumen microbial genes contribute little. *Scientific Reports* 8, 1–7.
- Meale, S.J., Li, S., Azevedo, P., Derakhshani, H., DeVries, T., Plaizier, J., Steele, M., Khafipour, E., 2017a. Weaning age influences the severity of gastrointestinal microbiome shifts in dairy calves. *Scientific Reports* 7, 1–13.
- Meale, S.J., Li, S., Azevedo, P., Derakhshani, H., Plaizier, J.C., Khafipour, E., Steele, M.A., 2016. Development of ruminal and fecal microbiomes are affected by weaning but not weaning strategy in dairy calves. *Frontiers in Microbiology* 7, 582.
- Meale, S.J., Morgavi, D.P., Cassar-Malek, I., Andueza, D., Ortigues-Marty, I., Robins, R. J., Schiphorst, A.-M., Laverroux, S., Graulet, B., Boudra, H., 2017b. Exploration of biological markers of feed efficiency in young bulls. *Journal of Agricultural and Food Chemistry* 65, 9817–9827.
- Mercadante, M.E.Z., Caliman, A.P.D.M., Canesin, R.C., Bonilha, S.F.M., Berndt, A., Frighetto, R.T.S., Magnani, E., Branco, R.H., 2015. Relationship between residual feed intake and enteric methane emission in Nelore cattle. *Revista Brasileira de Zootecnia* 44, 255–262.
- Muetzel, S., 2015. Framework & Milestone report—H<sub>2</sub> management in the rumen Final Report, MPI Technical Paper No: 2018/30. Ministry for Primary Industries, New Zealand Government, Wellington, New Zealand.
- Muetzel, S., Clark, H., 2015. Methane emissions from sheep fed fresh pasture. *New Zealand Journal of Agricultural Research* 58, 472–489.
- Nasrollahi, S.M., Meale, S.J., Morgavi, D.P., Schiphorst, A.M., Robins, R.J., Ortigues-Marty, I., Cantalapiedra-Hijar, G., 2020. The origin of N isotopic discrimination and its relationship with feed efficiency in fattening yearling bulls is diet-dependent. *PLOS One* 15, e0234344.
- Negussie, E., de Haas, Y., Dehareng, F., Dewhurst, R.J., Dijkstra, J., Gengler, N., Morgavi, D., Soyeurt, H., van Gastelen, S., Yan, T., 2017. Invited review: Large-scale indirect measurements for enteric methane emissions in dairy cattle: A review of proxies and their potential for use in management and breeding decisions. *Journal of Dairy Science* 100, 2433–2453.
- Neill, A.R., Grime, D.W., Dawson, R., 1978. Conversion of choline methyl groups through trimethylamine into methane in the rumen. *Biochemical Journal* 170, 529–535.
- Nkrumah, J., Basarab, J., Price, M., Okine, E., Ammoura, A., Guercio, S., Hansen, C., Li, C., Benkel, B., Murdoch, B., 2004. Different measures of energetic efficiency and their phenotypic relationships with growth, feed intake, and ultrasound and carcass merit in hybrid cattle. *Journal of Animal Science* 82, 2451–2459.
- Nkrumah, J., Okine, E., Mathison, G., Schmid, K., Li, C., Basarab, J., Price, M., Wang, Z., Moore, S., 2006. Relationships of feedlot feed efficiency, performance, and feeding behavior with metabolic rate, methane production, and energy partitioning in beef cattle. *Journal of Animal Science* 84, 145–153.
- Raisbeck, M., Riker, S., Tate, C., Jackson, R., Smith, M., Reddy, K., Zygmunt, J., 2008. Water quality for Wyoming livestock & wildlife: A review of the literature pertaining to health effects of inorganic contaminants, B-1183. University of Wyoming, Cheyenne, WY, USA.
- Rey, M., Enjalbert, F., Combes, S., Cauquil, L., Bouchez, O., Monteils, V., 2014. Establishment of ruminal bacterial community in dairy calves from birth to weaning is sequential. *Journal of Applied Microbiology* 116, 245–257.
- Richardson, E., Herd, R., 2004. Biological basis for variation in residual feed intake in beef cattle. 2. Synthesis of results following divergent selection. *Australian Journal of Experimental Agriculture* 44, 431–440.
- Rieu, F., Fonty, G., Gaillard, B., Gouet, P., 1990. Electron microscopy study of the bacteria adherent to the rumen wall in young conventional lambs. *Canadian Journal of Microbiology* 36, 140–144.
- Roehe, R., Dewhurst, R.J., Duthie, C.A., Rooke, J.A., McKain, N., Ross, D.W., Hyslop, J.J., Waterhouse, A., Freeman, T.C., Watson, M., 2016. Bovine host genetic variation influences rumen microbial methane production with best selection criterion for low methane emitting and efficiently feed converting hosts based on metagenomic gene abundance. *PLOS Genetics* 12, e1005846. <https://doi.org/10.1371/journal.pgen.1005846>.
- Roman, R., Wilcox, C., Martin, F., 2000. Estimates of repeatability and heritability of productive and reproductive traits in a herd of Jersey cattle. *Genetics and Molecular Biology* 23, 113–119.
- Sahoo, A., Kamra, D., Pathak, N., 2005. Pre- and postweaning attributes in faunated and ciliate-free calves fed calf starter with or without fish meal. *Journal of Dairy Science* 88, 2027–2036.
- Sander, E., Warner, R., Harrison, H., Loosli, J., 1959. The stimulatory effect of sodium butyrate and sodium propionate on the development of rumen mucosa in the young calf. *Journal of Dairy Science* 42, 1600–1605.
- Schenkel, F., Miller, S., Wilton, J., 2004. Genetic parameters and breed differences for feed efficiency, growth, and body composition traits of young beef bulls. *Canadian Journal of Animal Science* 84, 177–185.
- Schwarm, A., Schweigel-Röntgen, M., Kreuzer, M., Ortman, S., Gill, F., Kuhla, B., Meyer, U., Lohölter, M., Derno, M., 2015. Methane emission, digestive characteristics and faecal archaeol in heifers fed diets based on silage from brown midrib maize as compared to conventional maize. *Archives of Animal Nutrition* 69, 159–176.
- Seshadri, R., Leahy, S.C., Attwood, G.T., Teh, K.H., Lambie, S.C., Cookson, A.L., Eloe-Fadros, E.A., Pavlopoulos, G.A., Hadjithomas, M., Varghese, N.J., 2018. Cultivation and sequencing of rumen microbiome members from the Hungate1000 Collection. *Nature Biotechnology* 36, 359.
- Shingfield, K., 2000. Estimation of microbial protein supply in ruminant animals based on renal and mammary purine metabolite excretion. A review. *Journal of Animal and Feed Sciences* 9, 169–212.
- Soberon, F., Van Amburgh, M., 2013. Lactation Biology Symposium: The effect of nutrient intake from milk or milk replacer of preweaned dairy calves on lactation milk yield as adults: a meta-analysis of current data. *Journal of Animal Science* 91, 706–712.
- Steele, M.A., Penner, G.B., Chaucheyras-Durand, F., 2016. Development and physiology of the rumen and the lower gut: Targets for improving gut health. *Journal of Dairy Science* 99, 4955–4966.
- Stobo, I., Roy, J., Gaston, H.J., 1966. Rumen development in the calf: 1. The effect of diets containing different proportions of concentrates to hay on rumen development. *British Journal of Nutrition* 20, 171–188.
- Suárez, B., Van Reenen, C., Beldman, G., Van Delen, J., Dijkstra, J., Gerrits, W., 2006. Effects of supplementing concentrates differing in carbohydrate composition in veal calf diets: I. Animal performance and rumen fermentation characteristics. *Journal of Dairy Science* 89, 4365–4375.
- Talebi, M., 2012. Feed intake, feed efficiency, growth and their relationship with Kleiber ratio in Lori-Bakhtiari lambs. *Archiva Zootechnica* 15, 33.
- Tamate, H., McGilliard, A., Jacobson, N., Getty, R., 1962. Effect of various diets on the anatomical development of the stomach in the calf. *Journal of Dairy Science* 45, 408–420.
- Tamminga, S., 1979. Protein degradation in the forestomachs of ruminants. *Journal of Animal Science* 49, 1615–1630.
- Tedeschi, L., Fox, D., Russell, J., 2000. Accounting for ruminal deficiencies of nitrogen and branched-chain amino acids in the structure of the Cornell net carbohydrate and protein system. In: *Proceedings of the Cornell Nutrition Conference for Feed Manufacturers*, 24–26 October 2000, Rochester, New York, NY, USA, pp. 224–238. Article number 19.
- Van den Borne, J., Verstegen, M., Alferink, S., Giebels, R., Gerrits, W., 2006. Effects of feeding frequency and feeding level on nutrient utilization in heavy preruminant calves. *Journal of Dairy Science* 89, 3578–3586.

- Van Nevel, C., Demeyer, D., 1996. Control of rumen methanogenesis. *Environmental Monitoring and Assessment* 42, 73–97.
- Van Soest, P.J., 1996. Allometry and ecology of feeding behavior and digestive capacity in herbivores: a review. *Zoo Biology*: Published in affiliation with the American Zoo and Aquarium Association 15, 455–479.
- Vlaeminck, B., Dufour, C., Van Vuuren, A., Cabrita, A., Dewhurst, R.J., Demeyer, D., Fievez, V., 2005. Use of odd and branched-chain fatty acids in rumen contents and milk as a potential microbial marker. *Journal of Dairy Science* 88, 1031–1042.
- Vlaeminck, B., Fievez, V., Cabrita, A., Fonseca, A., Dewhurst, R., 2006a. Factors affecting odd-and branched-chain fatty acids in milk: A review. *Animal Feed Science and Technology* 131, 389–417.
- Vlaeminck, B., Fievez, V., Demeyer, D., Dewhurst, R.J., 2006b. Effect of forage: concentrate ratio on fatty acid composition of rumen bacteria isolated from ruminal and duodenal digesta. *Journal of Dairy Science* 89, 2668–2678.
- Waghorn, G., Hegarty, R., 2011. Lowering ruminant methane emissions through improved feed conversion efficiency. *Animal Feed Science and Technology* 166, 291–301.
- Wallace, R.J., Sasson, G., Garnsworthy, P.C., Tapio, I., Gregson, E., Bani, P., Huhtanen, P., Bayat, A.R., Strozzi, F., Biscarini, F., 2019. A heritable subset of the core rumen microbiome dictates dairy cow productivity and emissions. *Science. Advances* 5, eaav8391.
- Wang, Z., Nkrumah, J.D., Li, C., Basarab, J.A., Goonewardene, L.A., Okine, E.K., Crews Jr, D.H., Moore, S.S., 2006. Test duration for growth, feed intake, and feed efficiency in beef cattle using the GrowSafe System. *Journal of Animal Science* 84, 2289–2298.
- Wang, Z., Elekwachi, C., Jiao, J., Wang, M., Tang, S., Zhou, C., Tan, Z., Forster, R.J., 2017. Changes in metabolically active bacterial community during rumen development, and their alteration by rhubarb root powder revealed by 16S rRNA amplicon sequencing. *Frontiers in Microbiology* 8, 159.
- Westreicher-Kristen, E., Castro-Montoya, J., Hasler, M., Susenbeth, A., 2020. Relationship of milk odd-and branched-chain fatty acids with urine parameters and ruminal microbial protein synthesis in dairy cows fed different proportions of maize silage and red clover silage. *Animals* 10, 316.
- Wheadon, N., McGee, M., Edwards, G., Dewhurst, R., 2014. Plasma nitrogen isotopic fractionation and feed efficiency in growing beef heifers. *British Journal of Nutrition* 111, 1705–1711.
- Yáñez-Ruiz, D.R., Abecia, L., Newbold, C.J., 2015. Manipulating rumen microbiome and fermentation through interventions during early life: a review. *Frontiers in Microbiology* 6, 1133.